

Earliest land plants created modern levels of atmospheric oxygen

Timothy M. Lenton^{a,1}, Tais W. Dahl^b, Stuart J. Daines^a, Benjamin J. W. Mills^{a,c}, Kazumi Ozaki^d, Matthew R. Saltzman^e, and Philipp Porada^f

^aEarth System Science, College of Life and Environmental Sciences, University of Exeter, Exeter EX4 4QE, United Kingdom; ^bNatural History Museum of Denmark, University of Copenhagen, DK-1350 Copenhagen, Denmark; ^cSchool of Earth and Environment, University of Leeds, Leeds LS2 9JT, United Kingdom; ^dSchool of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA 30332-0340; ^eSchool of Earth Sciences, Ohio State University, Columbus, OH 43214; and ^fDepartment of Environmental Science and Analytical Chemistry, Stockholm University, SE-114 18 Stockholm, Sweden

Edited by Mark H. Thieme, University of California at San Diego, La Jolla, CA, and approved July 11, 2016 (received for review March 23, 2016)

The progressive oxygenation of the Earth's atmosphere was pivotal to the evolution of life, but the puzzle of when and how atmospheric oxygen (O₂) first approached modern levels (~21%) remains unresolved. Redox proxy data indicate the deep oceans were oxygenated during 435–392 Ma, and the appearance of fossil charcoal indicates O₂ >15–17% by 420–400 Ma. However, existing models have failed to predict oxygenation at this time. Here we show that the earliest plants, which colonized the land surface from ~470 Ma onward, were responsible for this mid-Paleozoic oxygenation event, through greatly increasing global organic carbon burial—the net long-term source of O₂. We use a trait-based ecophysiological model to predict that cryptogamic vegetation cover could have achieved ~30% of today's global terrestrial net primary productivity by ~445 Ma. Data from modern bryophytes suggests this plentiful early plant material had a much higher molar C:P ratio (~2,000) than marine biomass (~100), such that a given weathering flux of phosphorus could support more organic carbon burial. Furthermore, recent experiments suggest that early plants selectively increased the flux of phosphorus (relative to alkalinity) weathered from rocks. Combining these effects in a model of long-term biogeochemical cycling, we reproduce a sustained +2‰ increase in the carbonate carbon isotope (δ¹³C) record by ~445 Ma, and predict a corresponding rise in O₂ to present levels by 420–400 Ma, consistent with geochemical data. This oxygen rise represents a permanent shift in regulatory regime to one where fire-mediated negative feedbacks stabilize high O₂ levels.

oxygen | Paleozoic | phosphorus | plants | weathering

After the well-defined “Great Oxidation Event” 2.45–2.32 Ga, the trajectory of atmospheric oxygen is deeply uncertain (1, 2). Many recent studies, reviewed in refs. 3–5 have argued for a Neoproterozoic oxygenation event (>550 Ma)—of uncertain cause—and have linked it to the rise of animals, but this has been questioned given a lack of change in iron speciation ocean redox proxy data (6). Some models predict pO₂ ~1 present atmospheric level (PAL) already in the Early Paleozoic (7, 8), but this is at odds with data for widespread ocean anoxia (6, 9). The COPSE model we adapt here (10) predicts Early Paleozoic pO₂ ~0.2–0.5 PAL consistent with redox proxy data but, like the other models (7, 8), it does not predict a rise in oxygen until the advent of forests starting ~385 Ma, and continuing until ~300 Ma, which is too late to explain marked changes in geochemical data that occur before ~390 Ma (Fig. 1). The first appearance of fossil charcoal in the Late Silurian (11) and its ongoing occurrence through the Devonian (12) (Table S1), albeit rare and at low concentrations, indicates O₂ > 15–17% (vol) of the atmosphere (13) (or O₂ > ~0.7 PAL assuming a constant N₂ reservoir) already by ~420–400 Ma. [Under ideal conditions of ultradry fuel and forced airflow, smoldering fires may be sustained at O₂ > 10%, but this is not believed to be possible under natural conditions (14).] The molybdenum isotope record (9) indicates a fundamental shift in the redox state of the deep ocean from widespread anoxia to widespread oxygenation sometime during 435–392 Ma (between the Early Silurian and the Middle

Devonian). This ocean oxygenation is also supported by a Silurian increase in the C/S ratio of shales (15), and a shift in iron speciation data sometime during 435–387 Ma (6).

The persistent oxygenation of the ocean and appearance of charcoal can be explained by a rise in atmospheric oxygen occurring by ~400 Ma; this could be due to a persistent increase in oxygen source—considered here—or a decrease in oxygen sink (16), leading to a reorganization of the Earth's surface redox balance at a higher steady-state level for atmospheric O₂. The major long-term source of oxygen to the atmosphere is the burial of organic carbon in sedimentary rocks (which represents the net flux of photosynthesis minus various pathways of respiration and oxidation). Increases in global organic carbon burial are recorded as positive shifts in the isotopic composition of carbonate rocks (δ¹³C). Consistent with a rise in oxygen, the carbon isotope record (17) (Fig. 1) indicates a fundamental shift in baseline from ≤0‰ before the Late Ordovician to ~2‰ from ~445 Ma onward. Though there are many subsequent δ¹³C fluctuations, including drops back to 0‰, for example, at ~400 Ma, the long-term mean δ¹³C remains ~2‰ throughout the rest of the Paleozoic, the Mesozoic, and the Early Cenozoic (17), indicating a sustained increase in global organic carbon burial. Such a permanent shift requires a unidirectional driver that kicked in during the mid-Paleozoic. The evolution of land plants is the obvious candidate, with the first nonvascular plants (ancestors of extant mosses, liverworts, and hornworts) colonizing the land in the Middle to Late Ordovician (~470–445 Ma), followed by the first vascular plants in the Silurian (~445–420 Ma) and Early Devonian (~420–390 Ma; Fig. 1) (18, 19).

Significance

The rise of atmospheric oxygen over Earth's history has received much recent interdisciplinary attention. However, the puzzle of when and how atmospheric oxygen reached modern levels remains unresolved. Many recent studies have argued for a major oxygenation event—of uncertain cause—in the Neoproterozoic Era >541 Ma, enabling the rise of animals. Previous modelling work has predicted a late Paleozoic oxygen rise (<380 Ma) due to the rise of forests. Here we show that neither scenario is correct. Instead, the earliest plants, which colonized the land from 470 Ma onward, first increased atmospheric oxygen to present levels by 400 Ma, and this instigated fire-mediated feedbacks that have stabilized high oxygen levels ever since, shaping subsequent evolution.

Author contributions: T.M.L. designed research; T.M.L. and P.P. performed research; T.M.L., S.J.D., B.J.W.M., K.O., and P.P. contributed new reagents/analytic tools; T.M.L., T.W.D., K.O., M.R.S., and P.P. analyzed data; and T.M.L. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

¹To whom correspondence should be addressed. Email: t.m.lenton@exeter.ac.uk.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1604787113/-DCSupplemental.




Fig. 2. Predicted Late Ordovician (445 Ma) NPP. Result from ecophysiological model of cryptogamic vegetation cover driven by simulated Late Ordovician (445 Ma) climate, atmospheric $\text{CO}_2 = 8$ PAL, and atmospheric $\text{O}_2 = 0.6$ PAL (14 vol%), with no ice sheet mask. Simulated global NPP = $18.7 \text{ GtC}\cdot\text{y}^{-1}$.

Results and Discussion

To test our hypothesis, we revised the COPSE biogeochemical model (10) to better capture the early rise of plants and examine under what conditions it could explain the geochemical data (persistent rise to $\delta^{13}\text{C} \sim 2\text{‰}$ and the appearance of charcoal). The original baseline model (10) predicts early Paleozoic $\text{O}_2 \sim 0.23$ PAL at a reference time of 445 Ma, supported by an organic carbon burial flux of $\sim 4 \times 10^{12} \text{ mol}\cdot\text{y}^{-1}$ (about half the present-day value) with $\delta^{13}\text{C} = 0.03\text{‰}$. In this stable state, oxidative weathering of ancient

organic carbon is correspondingly reduced and its sensitivity to changes in O_2 provides a key negative feedback stabilizing O_2 . Key assumptions going into altering the forcing of the model are the global extent and associated productivity of early plants, the C/P ratio of plant material that was buried, and their effect (if any) on phosphorus weathering. To help parameterize these factors we drew on a mixture of experiments, existing data, and more detailed spatial modeling.

We used a trait-based spatial model of cryptogamic vegetation (i.e., bryophyte and lichen) cover (29, 30) driven by Late Ordovician climate simulations (31) at different atmospheric CO_2 levels to predict the potential global net primary productivity (NPP) of the early plant biosphere (32). At atmospheric $\text{CO}_2 = 8$ PAL, consistent with Late Ordovician glaciations (20), predicted global NPP is $\sim 19 \text{ GtC}\cdot\text{y}^{-1}$ (GtC, billion metric tons of carbon) (Fig. 2), $\sim 30\%$ of today. Predicted NPP is sensitive to variations in CO_2 and climate (Fig. S1), ice sheet cover (Fig. S2), and O_2 (Table S2), but is consistently higher than the $4.3 \text{ GtC}\cdot\text{y}^{-1}$ (7% of today) estimated elsewhere (33). In the original COPSE model (10), predicted NPP only reaches $\sim 5\%$ of today's value in the Late Ordovician and Silurian, but when we assume a stronger Late Ordovician phase of land colonization by nonvascular plants (following ref. 20; *SI Materials and Methods*), then COPSE predicts global NPP 30–40% of today (Fig. 3A), consistent with the detailed spatial model. In COPSE, this advent of early land plants alone, with no assumed effect on weathering fluxes, and assumed C/P = 1,000, increases total organic carbon burial by $\sim 25\%$, $\delta^{13}\text{C}$ by 0.5‰ , and atmospheric O_2 by 0.11 PAL (Fig. 3, blue).

We undertook a literature review of molar C/P ratios in extant bryophytes (Table S3) to test whether C/P = 1,000 is a reasonable




Fig. 3. Predictions of mid-Paleozoic global carbon cycle change due to early plants from the updated COPSE model. (A) NPP. (B) Organic carbon burial (both terrestrial and marine-derived material). (C) Carbonate carbon isotope record ($\delta^{13}\text{C}$). (D) Atmospheric O_2 . Note that fossil charcoal 420–400 Ma indicates $\text{O}_2 > 0.66\text{--}0.77$ PAL. (Further results of the same model runs are in Figs. S3 and S5.) Black dashed line indicates original baseline model run; blue, early plant colonization (C/P = 1,000); cyan, early plant colonization + C/P = 2,000; magenta, early plant colonization + biotic effects on silicate weathering (C/P = 1,000); green, early plant colonization + C/P = 2,000 + biotic effects on silicate weathering; yellow, early plant colonization + biotic effects on silicate weathering + 50% increase in P weathering; red, early plant colonization + C/P = 2,000 + biotic effects on silicate weathering + 25% increase in P weathering; black, early plant colonization + C/P = 2,000 + biotic effects on silicate weathering + spikes of P weathering.

assumption for early plants, and this gives a range of C/P = 800–4,300 with a mean of C/P ~1,900. Furthermore, Early Devonian coaly shales indicate extensive peatlands 410–400 Ma and have C/N of 44–119 (28), comparable to that in modern peatlands where N/P and C/P ratios tend to increase with depth to C/P > 3,000 (34). Taken together, these data suggest that assuming C/P = 1,000 for early plants is conservative. If instead we assume that buried early plant matter had C/P = 2,000, then given their productivity, even with no effect on weathering fluxes, this increases global organic carbon burial by ~50%, $\delta^{13}\text{C}$ by 1.1‰, and atmospheric O_2 by 0.27 PAL (Fig. 3, cyan).

Early plants could also have had a significant effect on weathering fluxes (20), because they and their fungal mycorrhizal symbionts evolved means of accessing rock-bound nutrients, notably phosphorus. Experimental work (20) has shown that a modern nonvascular plant, the moss *Physcomitrella patens*, amplifies the weathering of Ca ions 1.4- to 3.6-fold and Mg ions 1.5- to 5.4-fold from silicate rocks (granite–andesite), and amplifies the weathering of phosphorus from granite ~24-fold (range 15–43; *Materials and Methods*). Subsequent experiments (21) with the modern liverwort *Marchantia paleacea* found a 2.5- to 7-fold amplification of Ca weathering and a 9- to 13-fold amplification of P weathering from basalt. Both studies thus indicate preferential weathering of P relative to Ca and Mg (and corresponding alkalinity). The presence of these rock-weathering capabilities in two early diverging lineages (mosses and liverworts) suggests it is an ancestral trait. It has been argued (21, 33) that such large measured local effects would not have scaled up to significant global effects, because of low global NPP (33) and a limited depth of influence in the soil (21). However, we estimate much higher global NPP (Fig. 2) and weathering potential (32). We also note that extensive shallow water phosphate deposits in the Late Ordovician (35) indicate a marked increase in phosphorus input to the ocean (20).

If we include in COPSE an effect of early plants on silicate weathering following ref. 20, assuming C/P = 1,000, this increases organic carbon burial by ~35%, $\delta^{13}\text{C}$ by 0.7‰, and O_2 by 0.18 PAL (Fig. 3, magenta). The effect on O_2 is constrained because atmospheric CO_2 and temperature are reduced (20) such that the silicate weathering flux (and associated phosphorus flux) continues to match the degassing flux of CO_2 (Fig. S3). However, increases in carbonate weathering (enhanced by plants) and oxidative weathering (due to the rise in O_2) increase the overall phosphorus weathering flux, roughly doubling the O_2 rise due to terrestrial production of high C/P material alone. Assuming that buried early plant matter had a higher C/P = 2,000 causes larger increases in total organic carbon burial ~60%, $\delta^{13}\text{C}$ + 1.2‰, and atmospheric O_2 + 0.35 PAL (Fig. 3, green).

However, to reproduce the observed $\delta^{13}\text{C}$ + 2‰ shift requires the inclusion of some selective weathering of phosphorus by early plants. Assuming that early plants caused a sustained 50% increase in phosphorus weathering relative to bulk rock dissolution, with C/P = 1,000, increases total organic carbon burial by ~95%, $\delta^{13}\text{C}$ by 2.2‰, and O_2 by 0.74 PAL (to 0.97 PAL at 417 Ma; Fig. 3, yellow). Assuming a sustained 25% increase in phosphorus weathering relative to bulk rock and C/P = 2,000 increases organic carbon burial by ~90%, $\delta^{13}\text{C}$ by 2.1‰, and O_2 by 0.67 PAL (Fig. 3, red). Alternatively, a series of P weathering spikes designed to reproduce the observed sequence of positive $\delta^{13}\text{C}$ excursions (Fig. 1), combined with C/P = 2,000, produces a series of spikes in organic carbon burial and a peak increase of O_2 of 0.72 PAL at 407 Ma (Fig. 3, black). We hypothesize that these assumed weathering spikes could reflect phases of plant colonization (20, 36) followed by the establishment of phosphorus recycling ecosystems (20). However, direct evidence linking a phase of land colonization to enhanced weathering and a positive $\delta^{13}\text{C}$ excursion has only thus far been established for the Silurian–Devonian boundary excursion (36). Therefore, alternative hypotheses for short-lived positive $\delta^{13}\text{C}$ excursions should also be considered.

Regarding the simulated long-term ~2‰ rise in $\delta^{13}\text{C}$, this is smaller than would be expected from standard application of the simplified formula: $\delta^{13}\text{C}(\text{ocean}) = \delta^{13}\text{C}(\text{river}) + f_{\text{org}} \cdot \epsilon$, where f_{org} is the fraction of carbon buried as organic matter, ϵ is the fractionation between carbonates and organic matter, and both ϵ and $\delta^{13}\text{C}(\text{river})$ are usually assumed to be constant. In our COPSE simulations there is a fully interactive isotope mass balance, and these terms are not constant. The approximate doubling of organic carbon burial (with roughly constant carbonate burial) represents an increase from $f_{\text{org}} = 0.18$ to $f_{\text{org}} = 0.31$. However, the increase in burial of isotopically light organic carbon is counteracted by an increase in the oxidative weathering of isotopically light organic carbon, which lowers the $\delta^{13}\text{C}$ of riverine input to the ocean from approximately -5‰ to approximately -7.5‰, which is in turn partially counteracted by an increase in fractionation between carbonates and organic matter from $\epsilon \sim 27$ to ~ 30 ‰, due to increasing O_2 (somewhat counteracted by declining CO_2).

Sensitivity analyses (*SI Materials and Methods*) indicate that our results are robust. Varying the uplift and degassing forcing of the model within plausible bounds only causes ± 0.08 PAL variation in O_2 about the initial state (Fig. S4), although it does cause the effect of the same early plant forcing scenario to range over +0.4 to 1.0 PAL O_2 (Table S4). Including an additional negative feedback on O_2 from increased marine organic C/P burial ratios under anoxic waters (37), increases its initial early Paleozoic level to 0.54 PAL and reduces the effect of the same biological forcing scenarios on O_2 by ~10–30%, giving a maximum increase of +0.63 PAL (Table S5). However, because the initial O_2 is now higher, the final O_2 is also higher in all cases, and even scenarios without selective weathering of phosphorus could explain the appearance of charcoal ($\text{O}_2 > \sim 0.7$ PAL).

Our model makes additional predictions that can be tested against geochemical data—notably, it predicts a decline in pyrite sulfur burial and associated drop in $\delta^{34}\text{S}$ and increases in seawater SO_4 concentration and C/S burial ratio with the rise of the earliest plants (Fig. S5). This finding is broadly consistent with the sulfur isotope ($\delta^{34}\text{S}$) record (38–40), which shows a marked decline through the Silurian–Early Devonian from ~30 to ~18‰, although available data also suggest an earlier Late Ordovician–Early Silurian rise from ~25 to ~30‰, which the present model does not capture. The model is consistent with proxy reconstructions of seawater SO_4 concentration, which suggest an Ordovician–Silurian rise from ~6 to ~10 mM (41), and with a Silurian increase in the molar C/S ratio of shales from ~5 to ~16 (15).

Other processes not yet included in the model warrant future consideration—for example, the effect of increasing atmospheric mass on climate (42) and the effect of weathering forcing scenarios on $\delta^7\text{Li}$ and $^{87}\text{Sr}/^{86}\text{Sr}$, which enable additional tests against data.

Conclusion

Our model can only reproduce Paleozoic geochemical data if the rise of the earliest land plants caused a major oxygenation event of the Earth's atmosphere and oceans by ~400 Ma. We attribute this mid-Paleozoic oxygenation event to a persistent global increase in organic carbon burial supported by the high C/P ratio of early land plant material, augmented by a plant-driven increase in P weathering flux relative to the weathering flux of alkalinity. The $\delta^{13}\text{C}$ record suggests this increase in organic carbon burial was essentially permanent, producing a new dynamically stable state for atmospheric O_2 . In this new steady state, oxidative weathering was increased (becoming less sensitive to variations in O_2) and new fire-mediated negative feedbacks on O_2 were instigated that have played a key role in stabilizing atmospheric O_2 concentration up to the present day (22, 43). For the earliest land plants to be responsible for such a major mid-Paleozoic oxygenation event requires that they were much more productive and globally

25. Gerrienne P, et al. (2011) A simple type of wood in two Early Devonian plants. *Science* 333(6044):837.
26. Labeuw L, Martone PT, Boucher Y, Case RJ (2015) Ancient origin of the biosynthesis of lignin precursors. *Biol Direct* 10(1):23.
27. Montañez IP (2016) A Late Paleozoic climate window of opportunity. *Proc Natl Acad Sci USA* 113(9):2334–2336.
28. Kennedy KL, et al. (2013) Lower Devonian coaly shales of northern New Brunswick, Canada: Plant accumulations in the early stages of Terrestrial colonization. *J Sediment Res* 83(12):1202–1215.
29. Porada P, Weber B, Elbert W, Poschl U, Kleidon A (2013) Estimating global carbon uptake by lichens and bryophytes with a process-based model. *Biogeosciences* 10(11):6989–7033.
30. Porada P, Weber B, Elbert W, Pöschl U, Kleidon A (2014) Estimating impacts of lichens and bryophytes on global biogeochemical cycles. *Global Biogeochem Cycle*, 10.1002/2013GB004705.
31. Pohl A, Donnadiu Y, Le Hir G, Buoncristiani JF, Vennin E (2014) Effect of the Ordovician paleogeography on the (in)stability of the climate. *Clim Past* 10(6):2053–2066.
32. Porada P, et al. (2016) High potential for weathering and climate effects of non-vascular vegetation in the Late Ordovician. *Nat Commun*, 10.1038/ncomms12113.
33. Edwards D, Cherns L, Raven JA (2015) Could land-based early photosynthesizing ecosystems have bioengineered the planet in mid-Palaeozoic times? *Palaeontology* 58(5):803–837.
34. Wang M, Moore TR, Talbot J, Richard PJH (2014) The cascade of C:N:P stoichiometry in an ombrotrophic peatland: from plants to peat. *Environ Res Lett* 9(2):024003.
35. Pope MC, Steffen JB (2003) Widespread, prolonged late Middle to Late Ordovician upwelling in North America: A proxy record of glaciation? *Geology* 31(1):63–66.
36. Malkowski K, Racki G (2009) A global biogeochemical perturbation across the Silurian–Devonian boundary: Ocean–continent–biosphere feedbacks. *Palaeogeogr Palaeoclimatol Palaeoecol* 276(1–4):244–254.
37. Van Cappellen P, Ingall ED (1996) Redox stabilization of the atmosphere and oceans by phosphorus-limited marine productivity. *Science* 271:493–496.
38. Prokoph A, Shields GA, Veizer J (2008) Compilation and time-series analysis of a marine carbonate $\delta^{18}O$, $\delta^{13}C$, $87Sr/86Sr$ and $\delta^{34}S$ database through Earth history. *Earth Sci Rev* 87(3–4):113–133.
39. Gill BC, Lyons TW, Saltzman MR (2007) Parallel, high-resolution carbon and sulfur isotope records of the evolving Paleozoic marine sulfur reservoir. *Palaeogeogr Palaeoclimatol Palaeoecol* 256(3–4):156–173.
40. Jones DS, Fike DA (2013) Dynamic sulfur and carbon cycling through the end-Ordovician extinction revealed by paired sulfate–pyrite $\delta^{34}S$. *Earth Planet Sci Lett* 363:144–155.
41. Algeo TJ, Luo GM, Song HY, Lyons TW, Canfield DE (2015) Reconstruction of secular variation in seawater sulfate concentrations. *Biogeosciences* 12(7):2131–2151.
42. Poulsen CJ, Tabor C, White JD (2015) Climate change. Long-term climate forcing by atmospheric oxygen concentrations. *Science* 348(6240):1238–1241.
43. Kump LR (1988) Terrestrial feedback in atmospheric oxygen regulation by fire and phosphorus. *Nature* 335:152–154.
44. Rimmer SM, Hawkins SJ, Scott AC, Cressler WL (2015) The rise of fire: Fossil charcoal in late Devonian marine shales as an indicator of expanding terrestrial ecosystems, fire, and atmospheric change. *Am J Sci* 315(8):713–733.
45. Glasspool IJ, Scott AC (2010) Phanerozoic concentrations of atmospheric oxygen reconstructed from sedimentary charcoal. *Nat Geosci* 3(9):627–630.
46. Diessel CFK (2010) The stratigraphic distribution of inertinite. *Int J Coal Geol* 81(4):251–268.
47. Glasspool IJ, Scott AC, Waltham D, Pronina N, Shao L (2015) The impact of fire on the Late Paleozoic Earth system. *Front Plant Sci* 6:756.
48. Edwards D, Axe L (2004) Anatomical evidence in the detection of the earliest wildfires. *Palaios* 19(2):113–128.
49. Glasspool IJ, Edwards D, Axe L (2006) Charcoal in the Early Devonian: A wildfire-derived Konservat-Lagerstätte. *Rev Palaeobot Palynol* 142(3–4):131–136.
50. Pflug HD, Prossl KF (1989) Palynology in gneiss—Results from the continental deep drilling program. *Naturwissenschaften* 76(12):565–567.
51. Pflug HD, Prossl KF (1991) Palynostratigraphical and paleobotanical studies in the pilot hole of the German continental deep drilling programme results and implications. *Sci Drill* 2(1):13–33.
52. Wollenweber J, et al. (2006) Characterisation of non-extractable macromolecular organic matter in Palaeozoic coals. *Palaeogeogr Palaeoclimatol Palaeoecol* 240(1–2):275–304.
53. Peppers RA, Damberger HH (1969) *Palynology and Petrography of a Middle Devonian Coal in Illinois* (Illinois State Geological Survey, Urbana, IL).
54. Ammosov II (1964) Petrographic composition of coals in the U. S. S. R. and some of its changes. *Int Geol Rev* 6(10):1798–1804.
55. Volkova IB (1994) Nature and composition of the Devonian coals of Russia. *Energy Fuels* 8(6):1489–1493.
56. Patrakov YF, Kamyanova VF, Fedyayeva ON (2005) A structural model of the organic matter of Barzass liptobliolitic coal. *Fuel* 84(2–3):189–199.
57. Sharypov VI, Kuznetsov BN, Beregovtsova NG, Startsev AN, Parmon VN (2006) Catalytic hydroliquefaction of Barzass liptobliolitic coal in a petroleum residue as a solvent. *Fuel* 85(7–8):918–922.
58. Ghori KAR (1999) *Silurian-Devonian Petroleum Source-Rock Potential and Thermal History, Carnarvon Basin, Western Australia* (Geological Survey of Western Australia, Perth, Australia).
59. Xu H-H, et al. (2012) Mid Devonian megaspores from Yunnan and North Xinjiang, China: Their palaeogeographical and palaeoenvironmental significances. *Palaeoworld* 21(1):11–19.
60. Yang Y, Zou R, Shi Z, Jiang R (1996) *Atlas for Coal Petrography of China* (China Univ of Mining and Technology Press, Beijing).
61. Dai S, Han D, Chou C-L (2006) Petrography and geochemistry of the Middle Devonian coal from Luquan, Yunnan Province, China. *Fuel* 85(4):456–464.
62. Goodarzi F, Gentsis T, Embry AF (1989) Organic petrology of two coal-bearing sequences from the Middle to Upper Devonian of Melville Island, Arctic Canada. Working paper (Geological Survey of Canada, Ottawa). Available at ftp.maps.canada.ca/pub/nrcan_rncan/publications/ess_sst/126/126737/pa_89_08.pdf.
63. Goodarzi F, Goodbody Q (1990) Nature and depositional environment of Devonian coals from western Melville Island, Arctic Canada. *Int J Coal Geol* 14(3):175–196.
64. Gentsis T, Goodarzi F (1991) Petrology, depositional environment and utilization potential of Devonian channel coals from Melville Island, Canadian Arctic Islands. *Bull Soc Geol Fr* 162(2):239–253.
65. Fowler MG, Goodarzi F, Gentsis T, Brooks PW (1991) Hydrocarbon potential of Middle and Upper Devonian coals from Melville Island, Arctic Canada. *Org Geochem* 17(6):681–694.
66. Michelsen JK, Khorasani GK (1991) A regional study on coals from Svalbard; Organic facies, maturity and thermal history. *Bull Soc Geol Fr* 162(2):385–397.
67. Rimmer SM, Thompson JA, Goodnight SA, Robl TL (2004) Multiple controls on the preservation of organic matter in Devonian–Mississippian marine black shales: Geochemical and petrographic evidence. *Palaeogeogr Palaeoclimatol Palaeoecol* 215(1–2):125–154.
68. Marynowski L, Filipiak P (2007) Water column euxinia and wildfire evidence during deposition of the Upper Famennian Hangenberg event horizon from the Holy Cross Mountains (central Poland). *Geol Mag* 144(03):569–595.
69. Cressler WL (2001) Evidence of earliest known wildfires. *Palaios* 16(2):171–174.
70. Rowe NP, Jones TP (2000) Devonian charcoal. *Palaeogeogr Palaeoclimatol Palaeoecol* 164(1–4):331–338.
71. Fairon-Demaret M, Hartkopf-Fröder C (2004) Late Famennian plant mesofossils from the Refrath 1 Borehole (Bergisch Gladbach-Paffrath Syncline; Ardennes-Rhenish Massif, Germany). *CFS Courier Forschungsinstitut Senckenberg* 251:89–121.
72. Prestianni C, Decombeix A-L, Thorez J, Fokan D, Gerrienne P (2010) Famennian charcoal of Belgium. *Palaeogeogr Palaeoclimatol Palaeoecol* 291(1–2):60–71.
73. Zhou L, et al. (2011) A new paleoenvironmental index for anoxic events—Mo isotopes in black shales from Upper Yangtze marine sediments. *Sci China Earth Sci* 54(7):1024–1033.
74. Herrmann AD, et al. (2012) Anomalous molybdenum isotope trends in Upper Pennsylvanian euxinic facies: Significance for use of $\delta^{98}Mo$ as a global marine redox proxy. *Chem Geol* 324–325:87–98.
75. Goldberg T, et al. (2013) Resolution of inter-laboratory discrepancies in Mo isotope data: An intercalibration. *J Anal At Spectrom* 28(5):724–735.
76. Nägler TF, et al. (2014) Proposal for an international molybdenum isotope measurement standard and data representation. *Geostand Geoanal Res* 38(2):149–151.
77. Canfield DE, Poulton SW, Narbonne GM (2007) Late-Neoproterozoic deep-ocean oxygenation and the rise of animal life. *Science* 315(5808):92–95.
78. Scott C, Lyons TW (2012) Contrasting molybdenum cycling and isotopic properties in euxinic versus non-euxinic sediments and sedimentary rocks: Refining the paleoproxies. *Chem Geol* 324–325:19–27.
79. Dahl TW, et al. (2013) Tracing euxinia by molybdenum concentrations in sediments using handheld X-ray fluorescence spectroscopy (HHXRF). *Chem Geol* 360–361:241–251.
80. Chapin FS, Johnson DA, McKendrick JD (1980) Seasonal movement of nutrients in plants of differing growth form in an Alaskan tundra ecosystem: Implications for herbivory. *J Ecol* 68(1):189–209.
81. Chapin FS (1989) The cost of tundra plant structures: Evaluation of concepts and currencies. *Am Nat* 133(1):1–19.
82. Chapin FS, Shaver GR (1989) Differences in growth and nutrient use among Arctic plant growth forms. *Funct Ecol* 3(1):73–80.
83. Shaver GR, Chapin FS (1991) Production: Biomass relationships and element cycling in contrasting Arctic vegetation types. *Ecol Monogr* 61(1):1–31.
84. Aerts R, Verhoeven JTA, Whigham DF (1999) Plant-mediated controls on nutrient cycling in temperate fens and bogs. *Ecology* 80(7):2170–2181.
85. Riis T, Olesen B, Katborg CK, Christoffersen KS (2010) Growth rate of an aquatic bryophyte (*Warnstorfia fluitans* (Hedw.) Loeske) from a high Arctic lake: Effect of nutrient concentration. *Arctic* 63(1):100–106.
86. Waite M, Sack L (2011) Does global stoichiometric theory apply to bryophytes? Tests across an elevation \times soil age ecosystem matrix on Mauna Loa, Hawaii. *J Ecol* 99(1):122–134.
87. Wang M, Moore T (2014) Carbon, nitrogen, phosphorus, and potassium stoichiometry in an ombrotrophic peatland reflects plant functional type. *Ecosystems (N Y)* 17(4):673–684.
88. Larmola T, et al. (2014) Methanotrophy induces nitrogen fixation during peatland development. *Proc Natl Acad Sci USA* 111(2):734–739.
89. Delgado V, Eddera A, Santamaría JM (2013) Nitrogen and carbon contents and $\delta^{15}N$ and $\delta^{13}C$ signatures in six bryophyte species: Assessment of long-term deposition changes (1980–2010) in Spanish beech forests. *Glob Change Biol* 19(7):2221–2228.
90. Elbert W, et al. (2012) Contribution of cryptogamic covers to the global cycles of carbon and nitrogen. *Nat Geosci* 5(7):459–462.
91. Royer DL, Donnadiu Y, Park J, Kowalczyk J, Goddérís Y (2014) Error analysis of CO₂ and O₂ estimates from the long-term geochemical model GEOCARBSULF. *Am J Sci* 314(9):1259–1283.